arXiv:math/0303328v1 [math.GT] 26 Mar 2003

(-2,3,7)-PRETZEL KNOT AND REEBLESS FOLIATION

JINHA JUN

communicated by Rachel Roberts

Abstract. If $p/q > 18$, p is odd, and $p/q \neq 37/2$, (p,q)-Dehn surgery for the (-2,3,7)-pretzel knot produces a 3-manifold without Reebless foliation.

1. INTRODUCTION

Every closed orientable 3-manifold admits a foliation with Reeb components [Rol]. On the contrary, Reebless foliation $\mathcal{F}$ reflects the topological information of the ambient manifold $M \supset \mathcal{F}$. Novikov[No] showed that leaves of $\mathcal{F}$ are π_1-injective and $\pi_2(M) = 0$. Rosenberg[Ros] showed M is irreducible or $M \approx S^2 \times S^1$. It follows that $\mathcal{F}$ lifts to $\widetilde{\mathcal{F}}$ which has planar leaves in the universal cover $\widetilde{M}$. Palmeira[Pa] proved that any simply connected $(n+1)$-manifold, $n \geq 2$, admitting a smooth foliations by planar leaves with codim=1 is diffeomorphic to $\mathbb{R}^{n+1}$. It follows that the universal cover of M is homeomorphic to $\mathbb{R}^3$(see also [CC] for the proof). Especially, M is irreducible and $\pi_1(M)$ is infinite.



Figure 1. (-2,3,7)-pretzel knot

Let E_K be the (-2,3,7)-pretzel knot complement (Fig 1). And let $E_K(p/q)$ denote the 3-manifold obtained by (p,q)-Dehn surgery along the (-2,3,7)-pretzel knot. It is known that there is no closed essential surface in E_K and the boundary slopes are 0/1, 16/1, 37/2, and 20/1 [HO][Oe]. Furthermore, $E_K(16/1), E_K(37/2)$, and $E_K(20/1)$ are toroidal [HO]. E_K admits (finite) cyclic surgery along 18/1 and 19/1[FS]. And Blieler and Hodgson[BH] showed $E_K(17/1)$ is a Seifert fibered space with finite fundamental group. In particular, $E_K(17/1), E_K(18/1)$, and $E_K(19/1)$

2000 *Mathematics Subject Classification.* Primary 57M25; Secondary 57R30.

Key words and phrases. (-2,3,7)-pretzel knot, Reebless foliation, essential lamination, Dehn surgery, group action.

The author was partially supported by BK21.

have no Reebless foliation by virtue of Rosenberg's Theorem. E_K is fibered over the circle with genus 5 surface whose monodromy is pseudo-Anosov and hence hyperbolic. The suspension of the stable laminations gives an essential lamination $\mathcal{L}$ in E_K with degeneracy slope =1(18/1) [Ga]. $\mathcal{L}$ remains essential in $E_K(p/q)$ if $|p-18q|>1$ [GO]. If p is even, $\mathcal{L}$ extends to a taut foliation in $E_K(p/q)$ by filling complementary regions with a bundle of monkey saddle except $p/q=18/1$.

Using the technique in [Rob] and [Li], one can prove that $E_K(p/q)$ has a Reebless foliation if $p/q \in (-\infty, 9)$. This is done by attaching product disks to the fibers.

Theorem 1.1 (**Main Theorem**). *If $p/q > 18$, p is odd, and $p/q \neq 37/2$, then $E_K(p/q)$ does not admit a Reebless foliation.*

The quotient space $\widetilde{M}/\widetilde{\mathcal{F}}$ is called the leaf space. An open transversal to leaves gives an 1-manifold structure if $\mathcal{F}$ is a Reebless foliation. The leaf space is a non-Hausdorff simply connected 1-manifold. There is a natural action of $\pi_1(M)$ on the leaf space induced from the action on $\widetilde{M}$. And this action has no global fixed point (see [Pa]). We will prove the Main theorem by showing there is no nontrivial π_1-action on any leaf space. Our technique are much the same as in [RRS].

Calegari and Dunfield [CD] notice that $\mathcal{F}$ gives rise to a faithful π_1-action on the a (universal) circle. They showed there is no taut foliation in the Weeks manifold (the closed hyperbolic 3-manifold with smallest known volume).

Our method is not applied to the case $p/q = 37/2$. Indeed, $E_K(37/2)$ contains a Reebless foliation. But this foliation is not taut, because it has dead-end components. The following is commented by Rachel Roberts.

Lemma 1.2. *$E_K(37/2)$ does contain a Reebless foliation.*

Proof: Eudave-Moñoz[Eu] showed that $E_K(37/2)$ is decomposed along the incompressible torus T^2 into $E_K(37/2) = X_L \cup_{T^2} X_R$, where X_L (respectively, X_R) is the left-handed (respectively, right-handed) trefoil knot complement.

Since X_L (respectively, X_R) is fibered, take the leaves of the foliations which meet $\partial X_L = T^2 = \partial X_R$ in simple closed curves of longitudinal slope and spiral them in a neighborhood of the torus. By adding T^2 as a leaf, neither side is a solid torus and so the resulting is Reebless. □

It is remarkable that any essential lamination in $E_K(37/2)$ contains torus T^2 as a leaf [BNR]. The proof of Main Theorem also can be used to show there is no transversely oriented essential lamination except $p/q = 37/2$. The following theorem immediately follows from the results of [RRS].

Theorem 1.3. *If $p/q > 18$, p is odd, and $p/q \neq 37/2$, (p/q)-Dehn surgery for the (-2,3,7)-pretzel knot gives a 3-manifold without transversely oriented essential lamination.*

If M contains an essential lamination with no isolated leaf, a leaf space corresponds to $\mathbb{R}$-order tree [GO]. In this case, π_1 acts on $\mathbb{R}$-order tree instead.

Since $E_K(37/2)$ is Haken, it contains transversely oriented essential lamination. In fact, there is the suspension of the stable lamination in $E_K(p/q)$ which remains essential when $|p-18q|>1$. Main Theorem and the argument above imply the following.

Corollary 1.4. *If $p/q > 19$ and p is odd, $E_K(p/q)$ contains essential lamination but does not admit any Reebless foliation.*

This paper is organized as follows. In Section 2, we discuss some basic properties of $\pi_1(E_K(p/q))$. Section 3 gives an outline of theory of group actions on (non-Hausdorff) simply connected 1-manifold. In Section 4, we will prove the nonexistence of $\mathbb{R}$-covered foliation in $E_K(p/q)$. And the proof of the Main Theorem is contained in Section 5 and 6.

All results in this paper were obtained while the author was visiting professor Rachel Roberts and professor John Shareshian in the Washington University in 2002. This paper would not be possible without their help. The author would like to express thanks to them for their hospitality that makes visit to St. Louis enjoyable and remarkable.

2. Fundamental Group

This section contains useful properties and a presentation of (-2,3,7)-pretzel knot group. In later sections, we will analyze the group actions on an orientable (non-Hausdorff) 1-manifold. The following proposition implies the action can be restricted to the orientation preserving one. Let J be a knot in S^3 and E_J be the exterior of J. Set $G_0 = \pi_1(E_J(p/q))$.

Proposition 2.1. *If p is odd, G_0 does not contain index 2 subgroup.*

Proof: Suppose contrary that there is a subgroup H with $[G_0 : H] = 2$. Since $G_0/H \cong \mathbf{Z}/2\mathbf{Z}$ is abelian, the commutator subgroup $[G_0, G_0]$ is a subgroup of H. Note that $G_0/[G_0, G_0] \cong H_1(E_J(p/q)) = \mathbf{Z}/p\mathbf{Z}$. Therefore we have a commutative diagram below. Because p is odd, we get a contradiction.

$$\begin{tikzcd}
G_0 \arrow[rr] \arrow[dr] & & G_0/[G_0,G_0] \cong \mathbf{Z}/p\mathbf{Z} \arrow[dl] \\
& G_0/H \cong \mathbf{Z}/2\mathbf{Z} &
\end{tikzcd}$$

□

Corollary 2.2. *Let X be any oriented manifold and let*

$$\Psi : G_0 \to Homeo(X)$$

be any homomorphism. Then $\Psi(G_0) \leq Homeo^+(X)$.

Proof: Suppose otherwise. Note that $[Homeo(X) : Homeo^+(X)] \leq 2$. Then $\Psi^{-1}(Homeo^+(X))$ is an index 2 subgroup of G_0. By Proposition 2.1, it is impossible. □

Using the computer program SNAPPEA [We], we can obtain a presentation of the fundamental group of the knot (or link) complement and the peripheral words using an ideal tetrahedra decomposition. Denote (-2,3,7)-pretzel knot complement by E_K. The fundamental group of E_K and the meridian m and longitude l are

$$\pi_1(E_K) = < a, b \mid a^2ba^2b^2a^{-1}b^2 >,$$
$$m = a^{-1}b^{-2}, \quad l = ab^{-1}a^2m^{-18}.$$

Of course, we have

$$a^2ba^2b^2a^{-1}b^2 = 1. \tag{2.1}$$

Let $G(p,q) := \pi_1(E_K(p/q)) =< a,b \mid a^2ba^2b^2a^{-1}b^2, m^pl^q > .$

Lemma 2.3 (Lemma 3.4, [RRS]). *There is some* $k \in G(p,q)$ *such that* $m = k^q$ *and* $l = k^{-p}$*.*

Now we have

$$l = ab^{-1}a^2m^{-18} \Longleftrightarrow k^{p-18q} = a^{-2}ba^{-1}. \tag{2.2}$$

The following relation plays a central role in our proof.

$$a^3k^{p-18q}a^3 = a^3(a^{-2}ba^{-1})a^3 = aba^2 = a^{-1}b^{-2}ab^{-2} = ma^2m. \tag{2.3}$$

3. Group action on the leaf space

We begin with a short exposition of the theory of group actions on non-Hausdorff simply connected 1-manifold, taken from [RRS]. Let $\mathcal{F}$ be a Reebless foliation in M. Then $\mathcal{F}$ can be lifted to $\widetilde{\mathcal{F}}$ in the universal cover $\widetilde{M}$. The quotient space $\mathcal{T} = \widetilde{M}/\widetilde{\mathcal{F}}$ is called the *leaf space*. The leaf space $\mathcal{T}$ is a simply connected, 2nd countable 1-manifold [CC]. But, in general, it is not necessarily Hausdorff. Moreover there is an 1-1 correspond between simply connected 1-manifolds and planar foliations in $\mathbb{R}^3$ up to conjugate by Palmeira [Pa] (see also [CC] for details). Gabai and Kazez [GK] extends this relation to the essential laminations and $\mathbb{R}$-order trees.

We recall here some terminology and definitions in [RRS]. Given $x, y \in \mathcal{T}$, we consider the *geodesic spine*

$$[[x,y]] = \{z \in \mathcal{T} | x, y \text{ lie in distinct components of } \mathcal{T} \setminus \{z\}\} \cup \{x,y\}$$

from x to y. $[[x,y]]$ is the union of a finite number of disjoint (possibly, degenerate) closed intervals.

$$[[x,y]] = [x,y_1] \cup [x_2,y_2] \cup \cdots \cup [x_n,y],$$

where y_i is not separated from x_{i+1}. Set

$$d(x,y) = n - 1.$$

Obviously, if $y \in [[x.z]]$ for some $x, y, z \in \mathcal{T}$

$$d(x,z) = d(x,y) + d(y,z).$$

Let us call a subset X of $\mathcal{T}$ *spine-connected* if for all $x, y \in X$, $[[x,y]] \subset X$.

Fix an orientation on $\mathcal{T}$. For $x \in \mathcal{T}$, $\mathcal{T} \setminus \{x\}$ has exactly two components since $\mathcal{T}$ is simply connected. If U is a connected Euclidean neighborhood of x, the two components of $U \setminus \{x\}$ lie in distinct components of $\mathcal{T} \setminus \{x\}$(Exercise C.1.4, [CC]). Only one component, say, U^+ is in the *positive* direction of x. Let x^+ be the component of $\mathcal{T} \setminus \{x\}$ containing U^+ and let x^- be the component $\mathcal{T} \setminus (x^+ \cup \{x\})$.

Now we define a partial relation $\leq$ on $\mathcal{T}$. For $x, y \in \mathcal{T}$,

$$x \leq y \Longleftrightarrow x^+ \supseteq y^+.$$

It follows that every map in $Homeo^+(\mathcal{T})$ preserves this order $\leq$.

Define a relation $\sim$ on $\mathcal{T}$ by $x \sim y$ if and only if x and y are not separated in $\mathcal{T}$. Set

$$[x] = \{y \in \mathcal{T} | y \sim x\}.$$

If $x \sim y$, let $\mathcal{T}_{\{x,y\}}$ denote the submanifold defined as follows:

- if $x \in y^+$ (equivalently, if $y \in x^+$), set $\mathcal{T}_{\{x,y\}} = \bigcap_{z \sim x \text{ and } z \sim y} z^+$, and
- if $x \in y^-$ (equivalently, if $y \in x^-$), set $\mathcal{T}_{\{x,y\}} = \bigcap_{z \sim x \text{ and } z \sim y} z^-$.

The relation $\sim$ is reflexive and symmetric, but not necessarily transitive. However, by Denjoy blowing up, we can modify $\mathcal{F}$ so that $\sim$ is an equivalent relation (see Appendix in [RRS]). In what follows we shall assume $\sim$ is an equivalence relation. Define the *Hausdorff tree* $\mathcal{T}_H = \mathcal{T}/\sim$.

If X, Y are disjoint, nonempty, spine-connected subsets of $\mathcal{T}$, the *bridge* from X to Y is the intersection of all paths in $\mathcal{T}$ with one end point in X and the other in Y. Similarly, we can define the bridge in $\mathcal{T}_H$.

For any group G acting on $\mathcal{T}$, if $g \in G$, denote

$$Fix(g) = \{x \in \mathcal{T} | xg = x\}$$

and

$$Nonsep(g) = \{x \in \mathcal{T} | xg \sim x\}.$$

We say the action is *trivial* or *has a global fixed point* if there is some $x \in \mathcal{T}$ such that $x \sim xg$ for all $g \in G$.

Define the *characteristic set* associated to g by

$$C_g = \{x \in \mathcal{T} | d(x, xg) \text{ is even }\}.$$

Lemma 3.1 (Lemma 4.7, [RRS]). *Let $x \in \mathcal{T}$. Then $x \in C_g$ if and only if x and xg are comparable with respect to the partial order $\leq$.*

Proposition 3.2 (Proposition 4.8, [RRS]). *Suppose $Nonsep(g) = \emptyset$. Then $C_g \neq \emptyset$ and for any $x \in C_g$,*

$$C_g = \bigcup_{n \in \mathbb{Z}} [[xg^n, xg^{n+1}]].$$

When $Nonsep(g) = \emptyset$, $A_g := C_g$ is called an *axis* for g. From Proposition 3.2, in $\mathcal{T}$, $A_g \approx \mathbb{R}$ or $A_g \approx \cup_{-\infty}^{\infty} [x_i, y_i]$, where $[x_i, y_i]$ is homeomorphic to a closed interval in $\mathbb{R}$, $[x_i, y_i] \cap [x_{i+1}, y_{i+1}] = \emptyset$ when $i \neq j$, $x_i \neq y_i$ and $y_i \sim x_{i+1}$ for all i, j. In each case, the action of g on A_g is conjugate to an action by translations. In $\mathcal{T}_H$, the image of A_g is homeomorphic to $\mathbb{R}$.

Suppose Y is a g-invariant embedded copy of $\mathbb{R}$ in $\mathcal{T}$ on which g acts freely. Then we call Y a *local axis* for g. Now suppose that $Nonsep(g) \neq \emptyset$ and let T_i for some $i \in \mathcal{I}$, denote the path components of $\mathcal{T} \setminus Nonsep(g)$. Notice that $T_i g = T_j$ for some $j \in \mathcal{I}$. Moreover, whenever $T_i g = T_i$, g acts freely on T_i, and hence this local action has an axis $A_g^i \subset T_i$. One can check that such an $A_g^i \approx \mathbb{R}$ and hence is an example of a local axis for g.

Lemma 3.3 (Lemma 4.10, [RRS]). *Suppose $Nonsep(g) \neq \emptyset$. Then*

$$C_g = Fix(g) \cup \{x \in \mathcal{T} | x \textit{ lies on a local axis for } g\}.$$

Lemma 3.4 (Corollary 4.12, [RRS]). *If there is some $x \in \mathcal{T}$ such that $d(x, xg) \neq 0$ is even, then $Nonsep(g) \neq \emptyset$.*

Lemma 3.5 (Corollary 4.13, [RRS]). *Let $g \in G$. Then both C_g and $C_g \cup Nonsep(g)$ are spine-connected.*

Sometimes it is useful to consider an object obtained by adding one points $\hat{x}$, called an *ideal point* of $\mathcal{T}$, to $\mathcal{T}$ for each $\sim$-equivalence class $[x]$ in $\mathcal{T}$ which contains more than one point. This object, denoted by $\widehat{\mathcal{T}}$, is called the *completion* of $\mathcal{T}$. We say that an ideal point $\hat{x}$ is a *source* if whenever y, z are distinct elements of $[x]$ we have $y \in z^-$ and we say that $\hat{x}$ is a *sink* if whenever y, z are distinct elements of $[x]$ we have $y \in z^+$. Note that every ideal point $\hat{x}$ is either a source or a sink. The action of any subgroup of $Homeo(\mathcal{T})$ extends to an action on $\widehat{\mathcal{T}}$ in the obvious way, that is, we set $\hat{x}g = \hat{y}$ if $[x]g = [y]$. We want to extend our partial order on $\mathcal{T}$ to $\widehat{\mathcal{T}}$ so that group actions on $\widehat{\mathcal{T}}$ obtained from orientation preserving actions on $\mathcal{T}$ preserve this extended partial order. For an ideal point $\hat{x}$, we define

$$\hat{x}^+ = \begin{cases} \bigcup_{y\in[x]}(\{y\} \cup y^+), & \hat{x} \text{ a source,} \\ \bigcap_{y\in[x]} y^+, & \hat{x} \text{ a sink,} \end{cases}$$

and set

$$\hat{x}^- = \mathcal{T} \setminus \hat{x}^+.$$

Note that $\hat{x}^+, \hat{x}^- \subset \mathcal{T}$. If $h \in Homeo^+(\mathcal{T})$, for $x, y \in \widehat{\mathcal{T}}$, we have $x^+ \subset y^+$ if and only if $(xh)^+ \subset (yh)^+$. So we extend the partial order $\leq$ in $\mathcal{T}$ to $\widehat{\mathcal{T}}$.

Whenever possible, we will use $\mathcal{T}_H$ instead of $\mathcal{T}$ to avoid tedious arguments when we deal with non-Hausdorff points and to use the simply connectedness.

Lemma 3.6 (Lemma 5.6, [RRS])**.** *Any nontrivial action of G on $\mathcal{T}$ canonically induces a nontrivial action of G on $\mathcal{T}_H$.*

The action of $\pi_1(M)$ on $\widetilde{M}$ induces a right action by homeomorphisms on $\mathcal{T}$. That is, there is a homomorphism

$$\Phi : \pi_1(M) \to Homeo(\mathcal{T}).$$

By Corollary 2.2, we can assume $\Phi : G(p,q) \to Homeo^+(\mathcal{T})$.

We set

$$\begin{aligned} \alpha &= \Phi(a) \\ \beta &= \Phi(b) \\ \mu &= \Phi(m) \\ \kappa &= \Phi(k). \end{aligned}$$

Thus we have

$$\alpha^2\beta\alpha^2\beta^2\alpha^{-1}\beta^2 = 1 \quad \text{by (2.1),} \tag{3.1}$$

$$\kappa^{p-18q} = \alpha^{-2}\beta\alpha^{-1} \quad \text{by (2.2),} \tag{3.2}$$

$$\alpha^3\kappa^{p-18q}\alpha^3 = \mu\alpha^2\mu \quad \text{by (2.3).} \tag{3.3}$$

By [CD, Theorem 7.9], we can assume Φ is injective, that is, π_1 acts faithfully. We will abuse the notation $G(p,q)$ for the image of Φ. When there is no ambiguity, we will simply say that $G(p,q)$, instead of $\Phi(G(p,q))$, acts on a leaf space $\mathcal{T}$.

4. ℝ-COVERED FOLIATION

In this section, we will prove nonexistence of ℝ-covered foliation in $E_K(p/q)$ for $p/q \geq 10$.

The following lemma will be used in several times in the proof of Main Theorem. If one wish to analyze other 3-manifold group following [RRS], this lemma seems to be a criterion in choosing a presentation of a group.

Lemma 4.1. *Let $G(p,q)$ act on a partially ordered set P. Suppose that $G(p,q)$ preserves order. If some $x \in P$ satisfies either of the conditions*

(1) *$x\kappa = x$ and $x, x\alpha$ are related in P, or*

(2) *$x\alpha = x$ and $x, x\kappa$ are related in P*
then x is fixed by every $g \in G(p,q)$.

Proof: Suppose $x\kappa = x$. Then we can assume $x < x\alpha$ because $G(p,q) =< \kappa, \alpha >$. Then

$$x = x\mu = x\alpha^{-1}\beta^{-2} < x\beta^{-2}. \tag{4.1}$$

But $x = x\kappa^{p-18q} = x\alpha^{-2}\beta\alpha^{-1} < x\beta\alpha^{-1}$. This implies

$$x\alpha < x\beta.$$

Since $x < x\alpha$, we have $x < x\beta < x\beta^2$. Contradiction to (4.1).

Similarly, if $x\alpha = x$ we may assume $x < x\kappa$. Then

$$x < x\mu = x\alpha^{-1}\beta^{-2} = x\beta^{-2}. \tag{4.2}$$

But $x < x\kappa^{p-18q} = x\alpha^{-2}\beta\alpha^{-1} = x\beta\alpha^{-1}$. This shows

$$x < x\beta.$$

Hence $x < x\beta^2$. Contradiction to (4.2). □

Lemma 4.2. *If $q > 0$ and $x\kappa > x$ for all $x \in \mathbb{R}$ then $x > x\beta$.*

Proof: Since $x\alpha^{-1}\beta^{-2} = x\mu = x\kappa^q > x$ for all $x \in \mathbb{R}$, we have

$$x\alpha^{-1} > x\beta^2. \tag{4.3}$$

Then

$$\begin{aligned} x &= x\alpha^2\beta\alpha^2\beta^2\alpha^{-1}\beta^2 && \text{by (3.1)} \\ &< x\alpha^2\beta\alpha^2\alpha^{-1}\alpha^{-1}\alpha^{-1} && \text{by (4.3)} \\ &= x\alpha^2\beta\alpha^{-1} \end{aligned}$$

and $x\alpha\beta^{-1} < x\alpha^2$. By replacing $x\alpha$ with x we get

$$x\alpha > x\beta^{-1}, \text{ and} \tag{4.4}$$

$$x\alpha^2 > x\beta^{-2}. \tag{4.5}$$

Thus

$$\begin{aligned} x &= x\alpha^2\beta\alpha^2\beta^2\alpha^{-1}\beta^2 && \text{by (3.1)} \\ &> x\beta^{-2}\beta\beta^{-2}\beta^2\alpha^{-1}\beta^2 && \text{by (4.5)} \\ &= x\beta^{-1}\alpha^{-1}\beta^{-2} \\ &> x\beta^{-1}\beta^2\beta^2 && \text{by (4.3)} \\ &= x\beta^3 \end{aligned}$$

for all $x \in \mathbb{R}$. This implies $x > x\beta$. □

Proposition 4.3. *If $p/q \geq 10$ and $\phi : G(p,q) \longrightarrow Homeo^+(\mathbb{R})$ is any homomorphism then there is some $x \in \mathbb{R}$ which is fixed by every element of $\phi(G(p,q))$.*

Proof: By Lemma 4.1, we may assume $x\kappa > x$ for all $x \in \mathbb{R}$. Then $x > x\beta$ for all $x \in \mathbb{R}$ by Lemma 4.2. Since

$$x = x\alpha^2\beta\alpha^2\beta^2\alpha^{-1}\beta^2 < x\alpha^2\alpha^2\alpha^{-1} = x\alpha^3,$$

Lemma 4.1 implies that $x\alpha > x$ for all $x \in \mathbb{R}$.

Since $x\mu > x$, $x > x\alpha\beta^2$ for all $x \in \mathbb{R}$. Therefore

$$x = x\alpha^2\beta\alpha(\alpha\beta^2)\alpha^{-1}\beta^2 < x\alpha^2\beta\alpha\alpha^{-1}\beta^2 = x\alpha^2\beta^3. \tag{4.6}$$

Now, we will prove the following.

$$x\alpha > x\mu^2 \text{ for all } x \in \mathbb{R}. \tag{4.7}$$

To see this, note that this is equivalent to $x\alpha^2\beta^2\alpha\beta^2 > x$.

$$\begin{aligned} x(\alpha^2\beta^2)\alpha\beta^2 &> x\beta^{-1}\alpha\beta^2 \quad \text{by (4.6)} \\ &> x\beta^{-1}\beta^{-1}\beta^2 \text{ by (4.4)} \\ &= x. \end{aligned}$$

From (4.7),

$$\begin{aligned} x\alpha > x\mu^2 &\Leftrightarrow x > x\mu^2\alpha^{-1} \\ &\Leftrightarrow x\beta^{-2} > x\mu^2\alpha^{-1}\beta^{-2} = x\mu^3. \end{aligned}$$

So we have

$$x\beta^{-2} > x\mu^3. \tag{4.8}$$

It follows that

$$\begin{aligned} x\kappa^{18q-p} = x\alpha\beta^{-1}\alpha^2 &> x\alpha\beta^{-1}\beta^{-3} \text{ by (4.6)} \\ &> x\mu^2\beta^{-4} \quad \text{by (4.7)} \\ &> x\mu^2\mu^6 \quad \text{by (4.8)} \\ &= x\mu^8 \\ &= x\kappa^{8q}. \end{aligned}$$

Since we assume $x\kappa > x$, $18q - p > 10q$. Hence $p/q < 10$. Contradiction to the hypothesis. □

We suspect that the Proposition is still true for $p/q \geq 9$. On the contrary, it is likely that the taut foliation for the coefficients $p/q \in (-\infty, 9)$ are $\mathbb{R}$-covered.

5. *Nonsep*$(\kappa) = \emptyset$

In this section, we will show the Main Theorem when $Nonsep(\kappa) = \emptyset$. From now, we will assume $p/q > 18 \Leftrightarrow p - 18q > 0$, unless specified otherwise.

Proposition 5.1. *Suppose $Nonsep(\kappa) = \emptyset$. Then the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: Consider the action on $\mathcal{T}_H$. There are 3 cases for $A_\kappa \cap A_\kappa\alpha$.

1. $A_\kappa \cap A_\kappa\alpha = A_\kappa$.
2. $A_\kappa \cap A_\kappa\alpha$ is a nonempty proper closed connected subset of A_κ.
3. $A_\kappa \cap A_\kappa\alpha = \emptyset$.

For case (1), $A_\kappa \approx \mathbb{R}$ is invariant under $\mathrm{Im}\Phi$ and hence there is a fixed point in $\mathcal{T}_H$ by Lemma 4.3. Thus, there is a global fixed point in $\mathcal{T}$ by Lemma 3.6.

Case (2) will be proved in Lemma 5.4 and case (3) in the Lemma below. □

Lemma 5.2. *Suppose $Nonsep(\kappa) = \emptyset$. If $A_\kappa \cap A_\kappa\alpha = \emptyset$, the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: The relation (3.3) applies to give

$$A_\kappa \alpha^2 \mu = A_\kappa \mu \alpha^2 \mu = A_\kappa \alpha^3 \kappa^{p-18q} \alpha^3.$$

We will compare the bridges from A_κ to $A_\kappa \alpha^2 \mu$ and $A_\kappa \alpha^3 \kappa^{p-18q} \alpha^3$ to find a contradiction.

In $\mathcal{T}_H$, we define a total order on A_κ by $x \preceq x\kappa$ for all $x \in A_\kappa$. Let $[r, s]$ be the bridge from A_κ to $A_\kappa \alpha$ in $\mathcal{T}_H$.

(1) $A_\kappa \cap A_\kappa \alpha^2 \neq \emptyset$.

Since the bridge from $A_\kappa \alpha$ to $A_\kappa \alpha^2$ is $[r\alpha, s\alpha]$, $s = r\alpha$. Let $[x, y] = A_\kappa \cap A_\kappa \alpha^2$. Possibly, x or y is not finite.

(a) $A_\kappa \cap A_\kappa \alpha^3 = \emptyset$.

Then the bridge from A_κ to $A_\kappa \alpha^3$ begins at r. Hence the bridge from $A_\kappa \alpha^3$ to $A_\kappa \alpha^3 \kappa^{p-18q} \alpha^3$ begins at $r\kappa^{p-18q}\alpha^3$. From the Fig 2, we see that the end point of the bridge from $A_\kappa \alpha$ to $A_\kappa \alpha^2$ is

$$r\alpha^2 = \begin{cases} r & \text{if } x \preceq r \preceq y, \\ x & \text{if } r \preceq x, \\ y & \text{if } y \preceq r, \end{cases}$$

because the bridge from $A_\kappa \alpha$ to $A_\kappa \alpha^2$ is $[r\alpha, r\alpha^2]$. On the other hand, the bridge from $A_\kappa \alpha^3$ to $A_\kappa \alpha^2 \mu$ begins at

$$r\kappa^{p-18q}\alpha^3 = \begin{cases} r\alpha & \text{if } x \preceq r \preceq y, \\ x\alpha & \text{if } r \preceq x, \\ y\alpha & \text{if } y \preceq r. \end{cases}$$

For all three cases, we have $r\kappa^{p-18q}\alpha^3 = r\alpha^3 \Leftrightarrow r\kappa^{p-18q} = r$. But $r \prec r\kappa^{p-18q}$. Contradiction.



$x \preceq r \preceq y$ $r \preceq x$ $y \preceq r$

FIGURE 2. $A_\kappa \cap A_\kappa \alpha^3 = \emptyset$

(b) $A_\kappa \cap A_\kappa \alpha^3 \neq \emptyset$.

It follows that $A_\kappa \alpha^3 \cap A_\kappa \alpha^3 \kappa^{p-18q} \alpha^3 \neq \emptyset$. Equivalently, $A_\kappa \alpha^3 \cap A_\kappa \alpha^2 \mu \neq \emptyset$. But $A_\kappa \alpha^2 \cap A_\kappa \alpha^3 = \emptyset$. Let $[z, w] = A_\kappa \cap A_\kappa \alpha^3$. Then $x \preceq y \prec z \preceq w$. Since the bridge from $A_\kappa \alpha^2$ to $A_\kappa \alpha^3$ is $[r\alpha^2, r\alpha^3]$, $y = r\alpha^2$ and $z = r\alpha^3$. That is, $z = y\alpha \in A_\kappa \alpha$. In particular, $z \notin A_\kappa$. Contradiction.

(2) $A_\kappa \cap A_\kappa \alpha^2 = \emptyset$

Then the bridge from A_κ to $A_\kappa \alpha^2$ is $[r, s\alpha]$ (Fig 3). So the bridge from A_κ to $A_\kappa \alpha^2 \mu$ is $[r\mu, s\alpha\mu]$(Fig 3).



FIGURE 3. bridge from A_κ to $A_\kappa\mu\alpha^2\mu$

(a) $A_\kappa \cap A_\kappa\alpha^3 \neq \emptyset$.
Since the bridge from $A_\kappa\alpha^2$ to $A_\kappa\alpha^3$ is $[r\alpha^2, s\alpha^2]$ and $A_\kappa \cap A_\kappa\alpha^2 = \emptyset$, $s\alpha = r\alpha^2 \Leftrightarrow s = r\alpha$. See Figures 3 and 4. Note that $[r, r\alpha] \cap [r\alpha, r\alpha^2] = \{r\alpha\}$. So $[r\alpha, r\alpha^2] \cap [r\alpha^2, r\alpha^3] = \{r\alpha^2\}$. But the bridge $[r\alpha^2, r\alpha^3]$ should contain $[r, r\alpha^2]$. Contradiction.



FIGURE 4. $A_\kappa \cap A_\kappa\alpha^2 = \emptyset$ and $A_\kappa \cap A_\kappa\alpha^3 \neq \emptyset$

(b) $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$.
The bridge from A_κ to $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$ is $[r, s\alpha^2\kappa^{p-18q}\alpha^3]$(Fig 5).


$s\alpha^2\kappa^{p-18q}$
$A_\kappa\alpha^3\kappa^{p-18q}$
$s\alpha^{-1}$
$A_\kappa = A_\kappa\kappa^{p-18q}$
$r\kappa^{p-18q}$
$A_\kappa\alpha^{-3}$
$r\alpha^{-3}$
α^3
$s\alpha^2\kappa^{p-18q}\alpha^3$
$A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$
$s\alpha^2$
$A_\kappa\alpha^3$
$r\kappa^{p-18q}\alpha^3$
A_κ
r

FIGURE 5. bridge from A_κ to $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$

Consequently, $[r\mu, s\alpha\mu] = [r, s\alpha^2\kappa^{p-18q}\alpha^3]$. But $r \neq r\mu = r\kappa^q$ in $A_\kappa \subset \mathcal{T}_H$. Contradiction.

$\square$

Owing to the following lemma, we can rule out the case that α acts on A_κ with fixed points in $\mathcal{T}_H$ when we prove Lemma 5.4

Lemma 5.3. *Suppose that in $\mathcal{T}$, we have $Nonsep(\alpha) \cap A_\kappa \neq \emptyset$. Then the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: If $x \in Fix(\alpha) \cap A_\kappa \neq \emptyset$, then $d(x, x\kappa)$ is necessarily even, and hence x and $x\kappa$ are comparable with respect to $\leq$ on $\mathcal{T}$. Then Lemma 4.1 applies. So we

may assume that $Fix(\alpha) \cap A_\kappa = \emptyset$, and choose $x \in Nonsep(\alpha) \cap A_\kappa$.
Consider $[x] \cap A_\kappa$. Either $[x] \cap A_\kappa = \{x\}$ or $[x] \cap A_\kappa = \{x, y\}$ for some $y \neq x$. In the first case, Lemma 4.1 applied to the ideal point determined by $[x]$ shows that the action of $G(p,q)$ on $\mathcal{T}$ is trivial. Now we assume the second case. Since $x \in A_\kappa$, we may assume $x\kappa \leq x$ (Fig 6). Let $d(x, x\kappa) = 2n > 0$.



FIGURE 6. $Nonsep(\alpha) \cap A_\kappa \neq \emptyset$

There are 5 subcases:

(1) $x = x\alpha^2$

$$\begin{aligned} x > x\kappa^{p-18q} &= x\alpha^{-2}\beta\alpha^{-1} = x\beta\alpha^{-1} \\ &\Leftrightarrow x\alpha > x\beta. \end{aligned}$$

But $x > x\mu = x\alpha^{-1}\beta^{-2} \Leftrightarrow x\beta^2 > x\alpha^{-1} = x\alpha$. Therefore,

$$\begin{aligned} & x\beta^2 > x\alpha > x\beta \\ & \Rightarrow x\beta > x \\ & \Rightarrow x\alpha > x\beta > x \\ & \Rightarrow x\alpha > x \\ & \Rightarrow x = x\alpha^2 > x\alpha > x. \end{aligned}$$

Contradiction.

(2) $y = x\alpha^2$
$x\beta = x\alpha^2\kappa^{p-18q}\alpha = y\kappa^{p-18q}\alpha > x$ and $x\beta \in (x\alpha)^-$ (Fig 7). Then $x\beta^2 = x\mu^{-1}\alpha^{-1} \in (y\alpha^{-1})^- = (x\alpha)^-$ and $x\beta^2 > x\beta$. We now see that $x\beta \in [[x, x\beta^2]]$. $d(x\beta, x\beta^2) = d(x, x\beta) = d(x, x\alpha^2\kappa^{p-18q}\alpha) = d(x\alpha^{-1}, y\kappa^{p-18q}) = d(y, y\kappa^{p-18q}) = 2n(p-18q)$, because $x\alpha^{-1} \neq x$.

Therefore,

$$\begin{aligned} 4n(p-18q) &= d(x, x\beta) + d(x\beta, x\beta^2) \\ &= d(x, x\beta^2) \\ &= d(x, x\mu^{-1}\alpha^{-1}) \\ &= d(x\alpha, x\mu^{-1}) \\ &= 2nq. \end{aligned}$$

That is, $p/q = 37/2$. However, we have assumed that $p/q \neq 37/2$.

(3) $y \neq x\alpha^2$ and $x = x\alpha^3$ (hence $x \neq x\alpha^2$)
$d(x, x\alpha^3\kappa^{p-18q}\alpha^3) = d(x, x\kappa^{p-18q}) = 2n(p-18q)$. On the other hand, $d(x, x\mu\alpha^2\mu) = 4nq$ (Fig 8). Hence $2n(p-18q) = 4nq \Leftrightarrow p = 20q$. But we assume p is odd.

(4) $x \neq x\alpha^2, x\alpha^3$ and $y = x\alpha^3$ (hence $y \neq x\alpha^2$)
$x\alpha^3\kappa^{p-18q}\alpha^3 = y\kappa^{p-18q}\alpha^3 \in (x\alpha^3)^- = y^-$, since $y\kappa^{p-18q} \in x^-$. But $x\mu\alpha^2\mu \in y^+$ (Fig 8).

(5) $x \neq x\alpha^2, x\alpha^3$ and $y \neq x\alpha^2, x\alpha^3$
$x\mu\alpha^2\mu \in x^-$, but $x\alpha^3\kappa^{p-18q}\alpha^3 \in x^+$ (Fig 8, 9).



FIGURE 7. $y = x\alpha^2$

□



FIGURE 8. $x\mu\alpha^2\mu$



FIGURE 9. $x\alpha^3\kappa^{p-18q}\alpha^3$

The following lemma, together with Lemma 5.2, will complete the proof of Proposition 5.1.

Lemma 5.4. *Suppose that $A_\kappa \cap A_\kappa\alpha$ is a nonempty proper closed connected subset of A_κ. Then the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: Suppose otherwise. In $\mathcal{T}_H$, $A_\kappa \approx \mathbb{R}$. Let $\preceq$ denote the total order on A_κ specified by $x \preceq x\kappa$ for all $x \in A_\kappa$. With respect to this order, let r (respectively, s) denote the lower bound (respectively, upper bound), if it exists, of $A_\kappa \cap A_\kappa\alpha$. Otherwise, set $r = -\infty$ (respectively, $s = \infty$). Note that at least one of r and s is finite because the intersection is a proper subset.
When $r \neq s$ let $\preceq_\alpha$ denote a total order on $A_\kappa\alpha$ such that $\preceq$ and $\preceq_\alpha$ agree on

$A_\kappa \cap A_\kappa\alpha$. Similarly, we define $\preceq_{\alpha^2}$ (respectively, $\preceq_{\alpha^3}$) on $A_\kappa\alpha^2$ (respectively, $A_\kappa\alpha^3$) to agree with $\preceq_\alpha$ (respectively, $\preceq_{\alpha^2}$) on $A_\kappa\alpha \cap A_\kappa\alpha^2$ (respectively, $A_\kappa\alpha^2 \cap A_\kappa\alpha^3$).

Let Y be an embedded copy of $\mathbb{R}$ in $\mathcal{T}_H$ with a total order. Then the homeomorphism

$$\alpha^r : Y \to Y\alpha^r$$

is order-preserving or order-reversing if some total order is defined in $Y\alpha^r$. Suppose that $\alpha : (A_\kappa, \preceq) \to (A_\kappa\alpha, \preceq_\alpha)$ is order-preserving. Then $r\alpha \preceq_\alpha s\alpha$ and $r\alpha^{-1} \preceq s\alpha^{-1}$. Since $r\alpha, s\alpha \in A_\kappa\alpha^2$, $r\alpha \preceq_{\alpha^2} s\alpha$. Therefore the map $\alpha : (A_\kappa\alpha, \preceq_\alpha) \to (A_\kappa\alpha^2, \preceq_{\alpha^2})$ is order preserving. In particular, $r\alpha^2 \preceq_{\alpha^2} s\alpha^2 \Leftrightarrow r\alpha^2 \preceq_{\alpha^3} s\alpha^2$. Since $(r\alpha^{-1})\alpha^3 = r\alpha^2 \preceq_{\alpha^3} s\alpha^2 = (s\alpha^{-1})\alpha^3$, the map $\alpha^3 : (A_\kappa, \prec) \to (A_\kappa\alpha^3, \prec_{\alpha^3})$ is order preserving.

If r and s are finite and either

$$[r\alpha, s\alpha] \subset [r, s] \text{ or } [r, s] \subset [r\alpha, s\alpha],$$

the intermediate value theorem implies $Nonsep(\alpha) \cap A_\kappa \neq \emptyset$. This contradicts to Lemma 5.3.

Accordingly, we have 3 cases by symmetry (Fig 10),
(I) $r \prec_\alpha r\alpha \prec_\alpha s \prec_\alpha s\alpha$.
(II) $r \prec_\alpha r\alpha = s \prec_\alpha s\alpha$.
(III) $r \preceq_\alpha s \prec_\alpha r\alpha \preceq_\alpha s\alpha$ (if $\preceq_\alpha$ is defined) $\Leftrightarrow [r, s] \cap [r\alpha, s\alpha] = \emptyset \Leftrightarrow A_\kappa \cap A_\kappa\alpha^2 = \emptyset$.



FIGURE 10. $A_\kappa, A_\kappa\alpha,$ and $A_\kappa\alpha^2$

For (I), note that

$$r\alpha \prec_{\alpha^2} r\alpha^2 \prec_{\alpha^2} s\alpha \prec_{\alpha^2} s\alpha^2.$$

If $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$, then $s \prec_{\alpha^2} r\alpha^2$. Then the bridge from $A_\kappa\alpha^3$ to $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$ is a translation of the bridge from A_κ to $A_\kappa\alpha^3\kappa^{p-18q}$ by α^3.

$$[s\kappa^{p-18q}, r\alpha^2\kappa^{p-18q}]\alpha^3 = [s\kappa^{p-18q}\alpha^3, r\alpha^2\kappa^{p-18q}\alpha^3] \text{(see Fig 11).}$$

On the other hand, the bridge from $A_\kappa\alpha^3$ to $A_\kappa\mu\alpha^2\mu$ is

(1) $[r\alpha^2, s]$ when $r\alpha\mu \prec s$,
(2) $[r\alpha^2, r\alpha\mu]$ when $s \prec r\alpha\mu$,



FIGURE 11. $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$

(3) $[r\alpha^2, x]$ when $s = r\alpha\mu$ for some x.
See Fig 12.



FIGURE 12. bridge from $A_\kappa\alpha^3$ to $A_\kappa\mu\alpha^2\mu$

Case (1) , $\begin{cases} s\kappa^{p-18q}\alpha^3 = r\alpha^2, \\ r\alpha^2\kappa^{p-18q}\alpha^3 = s. \end{cases}$ Hence both r and s are finite. By (3.2), the second relation is equivalent to $s = r\beta\alpha^2$. Substituting this in the first relation yields

$$\begin{aligned} r\alpha^2 &= r\beta\alpha^2\kappa^{p-18q}\alpha^3 = r\beta\alpha^2(\alpha^{-2}\beta\alpha^{-1})\alpha^3 \\ &\Leftrightarrow r = r\beta^2 = r\mu^{-1}\alpha^{-1} \\ &\Leftrightarrow r\alpha = r\mu^{-1}. \end{aligned}$$

But $r\mu^{-1} \prec r \prec r\alpha$. Contradiction.

Case (2), $\begin{cases} s\kappa^{p-18q}\alpha^3 = r\alpha^2, \\ r\alpha^2\kappa^{p-18q}\alpha^3 = r\alpha\mu. \end{cases}$ Hence both r and s are finite. We have

$$\begin{aligned} r\alpha\mu &= r\alpha^2(\alpha^{-2}\beta\alpha^{-1})\alpha^3 = r\beta\alpha^2 \\ &\Leftrightarrow r = r\beta\alpha^2\mu^{-1}\alpha^{-1} = r\beta\alpha^2\beta^2 = r\alpha^{-2}\beta^{-2}\alpha = r\alpha^{-1}\mu\alpha \qquad (5.1) \\ &\Leftrightarrow r\alpha^{-1} = r\alpha^{-1}\mu. \end{aligned}$$

But $r\alpha^{-1} \prec r\alpha^{-1}\mu$. Contradiction.

Case (3), $\begin{cases} s\kappa^{p-18q}\alpha^3 = r\alpha^2, \\ r\alpha^2\kappa^{p-18q}\alpha^3 = x, \\ s = r\alpha\mu. \end{cases}$ Then both r and s are finite. We have

$$\begin{aligned} r\alpha\mu\kappa^{p-18q}\alpha^3 &= r\alpha^2 \\ \Leftrightarrow r\alpha\kappa^{p-17q} &= r\alpha^{-1}. \end{aligned}$$

But $r\alpha^{-1} \prec r \prec r\alpha\kappa^{p-17q}$. Contradiction.

Now we may assume $A_\kappa \cap A_\kappa\alpha^3 \neq \emptyset$ (Fig 13). Then $r\alpha^2 \preceq_{\alpha^2} s$. Hence the intersection between A_κ and $A_\kappa\alpha^3\kappa^{p-18q}$ is $[r\alpha^2\kappa^{p-18q}, s\kappa^{p-18q}]$ in $\prec$-order in A_κ. Since $\alpha^3 : (A_\kappa, \prec) \to (A_\kappa\alpha^3, \prec_{\alpha^3})$ is an order-preserving map, the intersection between $A_\kappa\alpha^3$ and $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$ is $[r\alpha^2\kappa^{p-18q}\alpha^3, s\kappa^{p-18q}\alpha^3]$ in $\prec_{\alpha^3}$-order in $A_\kappa\alpha^3$. On the other hand, the intersection between $A_\kappa\alpha^3$ and $A_\kappa\alpha^2\mu$ is, in $\prec_{\alpha^3}$-order,



FIGURE 13. $A_\kappa \cap A_\kappa\alpha^3 \neq \emptyset$

(1) $[r\alpha^2, s]$ when $r\alpha\mu \prec r\alpha^2$,
(2) $[r\alpha\mu, s]$ when $r\alpha^2 \prec r\alpha\mu \prec s$,
(3) $[x, s]$ when $r\alpha^2 = r\alpha\mu$,
(4) $[s, y]$ when $s = r\alpha\mu$.

See Fig 14.



$r\alpha\mu \prec r\alpha^2$ $r\alpha^2 \prec r\alpha\mu \prec s$

$r\alpha^2 = r\alpha\mu$ $s = r\alpha\mu$

FIGURE 14. $A_\kappa\alpha^3 \cap A_\kappa\mu\alpha^2\mu$

Since r and s are not necessarily finite, we will show a contradiction even when one of r and s is not finite. Recall that at least one of r and s is finite.

Case (1), $[r\alpha^2, s] = [r\alpha^2\kappa^{p-18q}\alpha^3, s\kappa^{p-18q}\alpha^3]$.

- $r\alpha^2 = r\alpha^2\kappa^{p-18q}\alpha^3 \Leftrightarrow r\alpha^{-1} = r\alpha^2\kappa^{p-18q}$. But $r\alpha^{-1} \prec r \prec r\alpha^2\kappa^{p-18q}$. Contradiction.
- $s = s\kappa^{p-18q}\alpha^3$. Then $s \prec s\kappa^{p-18q} \Leftrightarrow s\alpha^3 \prec_{\alpha^3} s\kappa^{p-18q}\alpha^3 = s$. But $s \preceq_\alpha s\alpha \Leftrightarrow s\alpha^{-1} \preceq s \Leftrightarrow s\alpha^2 \preceq_{\alpha^3} s\alpha^3$. Since $s \prec_{\alpha^3} s\alpha^2 \prec_{\alpha^3} s\alpha^3$, we get a contradiction.

Case (2), $[r\alpha\mu, s] = [r\alpha^2\kappa^{p-18q}\alpha^3, s\kappa^{p-18q}\alpha^3]$.

- $r\alpha\mu = r\alpha^2\kappa^{p-18q}\alpha^3$. Contradiction as shown in case(2) of (I).
- $s = s\kappa^{p-18q}\alpha^3$. Contradiction as in Case (1).

Case (3), $[x, s] = [r\alpha^2\kappa^{p-18q}\alpha^3, s\kappa^{p-18q}\alpha^3]$ and $r\alpha^2 = r\alpha\mu$.

- $x = r\alpha^2\kappa^{p-18q}\alpha^3 \preceq_{\alpha^3} r\alpha^2 \Rightarrow r\alpha^2\kappa^{p-18q} \preceq r\alpha^{-1}$. But $r\alpha^{-1} \prec r\alpha^2 \prec r\alpha^2\kappa^{p-18q}$. Contradiction.
- $s = s\kappa^{p-18q}\alpha^3$. Contradiction as in Case (1).

Case (4), $[s, y] = [r\alpha^2\kappa^{p-18q}\alpha^3, s\kappa^{p-18q}\alpha^3]$ and $s = r\alpha\mu$. Then both r and s are finite. We have $r\alpha\mu = s = r\alpha^2\kappa^{p-18q}\alpha^3$. Contradiction as in Case (2) of (I).

For (II), note that $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$. And every arguments in this case reduce to the case (I) with $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$ and $s \prec s\mu = r\alpha\mu$ (Fig 15).



Figure 15. Case (II) $s = r\alpha$

For (III), note that $r\alpha \preceq_{\alpha^2} s\alpha \prec_{\alpha^2} r\alpha^2 \preceq_{\alpha^2} s\alpha^2$. In particular, $A_\kappa \cap A_\kappa\alpha^3 = \emptyset$ (Fig 16).

The bridge from $A_\kappa\alpha^3$ to $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$ is $[s\kappa^{p-18q}, r\alpha^2\kappa^{p-18q}]\alpha^3$. On the other hand, the bridge from $A_\kappa\alpha^3$ to $A_\kappa\alpha^2\mu = A_\kappa\mu\alpha^2\mu$ is $[r\alpha^2, r\alpha\mu]$. So we have

- $s\kappa^{p-18q}\alpha^3 = r\alpha^2$ and
- $r\alpha\mu = r\alpha^2\kappa^{p-18q}\alpha^3$.

Hence r and s are finite and we get a contradiction as shown in 5.1.

Now we may assume the map $\alpha : (A_\kappa, \prec) \to (A_\kappa\alpha, \prec_\alpha)$ is order reversing. Equivalently, $s\alpha \prec_\alpha r\alpha$.

If $[r, s] \cap [s\alpha, r\alpha] \neq \emptyset$, intermediated value theorem is applied to show $Nonsep(\alpha) \cap A_\kappa \neq \emptyset$. Hence we assume $[r, s] \cap [s\alpha, r\alpha] = \emptyset$. By symmetry, we can assume

$$r \prec_\alpha s \prec_\alpha s\alpha \prec_\alpha r\alpha \text{ (and hence } s \text{ is finite).}$$



Figure 16. Case (III)



FIGURE 17. order reversing

The bridge from $A_\kappa\alpha^3$ to $A_\kappa\alpha^3\kappa^{p-18q}\alpha^3$ is $[s\kappa^{p-18q}\alpha^3, s\alpha^2\kappa^{p-18q}\alpha^3]$(Fig 17). The bridge from $A_\kappa\alpha^3$ to $A_\kappa\alpha^2\mu$ is $[s\alpha^2, s\alpha\mu]$. So we have

$$\begin{aligned}
&s\alpha^2\kappa^{p-18q}\alpha^3 = s\alpha\mu \\
&\Leftrightarrow s\beta\alpha^2 = s\beta^{-2} \\
&\Leftrightarrow s\beta\alpha^2\beta^2 = s \\
&\Leftrightarrow s\alpha^{-1}\mu\alpha = s \quad \text{by (3.1)} \\
&\Leftrightarrow s\alpha^{-1}\mu = s\alpha^{-1}.
\end{aligned}$$

But $s\alpha^{-1} \prec s\alpha^{-1}\mu$. Contradiction. □

The line of reasoning used in this section shows that one actually has

Lemma 5.5. *Suppose Y is a κ-invariant embedded copy of $\mathbb{R}$ in $\mathcal{T}$ on which κ acts freely. If*

- $\emptyset \neq Y \cap Y\alpha \subset [r, s]$ *for some* $r, s \in Y$, *or*
- $Y \cap Y\alpha = \emptyset$ *and the bridge from* Y *to* $Y\alpha$ *has the form* $[[r, s]]$ *for some* $r \sim r' \in Y$, $s \sim s' \in Y\alpha$,

then the action $G(p, q)$ on $\mathcal{T}$ has a global fixed point.

6. $Nonsep(\kappa) \neq \emptyset$

We will complete the proof of Main Theorem by showing below that there is no nontrivial action on $\mathcal{T}$ when $Nonsep(\kappa) \neq \emptyset$

Lemma 6.1. *There is no $x \in \mathcal{T}$ which is nonseparated by κ and α.*

Lemma 6.2. *If $Fix(\kappa^{p-18q}) \cap Nonsep(\kappa) \cap C_\alpha \neq \emptyset$, then the action $G(p, q)$ on $\mathcal{T}$ is trivial.*

Proof: Let $x \in Fix(\kappa^{p-18q}) \cap Nonsep(\kappa) \cap C_\alpha$. By Lemma 3.1, x and $x\alpha$ are comparable. If $x\alpha = x$, $x = x\kappa^{p-18q} = x\alpha^{-2}\beta\alpha^{-1} = x\beta\alpha^{-1} \Leftrightarrow x = x\beta$. Because

$G(p,q) =< \alpha, \beta >$, instead we can assume that $x < x\alpha$. Then

$$\begin{aligned} x = x\kappa^{p-18q} &= x\alpha^{-2}\beta\alpha^{-1} < x\beta\alpha^{-1} \\ &\Rightarrow x\alpha < x\beta \\ &\Rightarrow x < x\beta \\ &\Rightarrow x < x\beta^2 = x\mu^{-1}\alpha^{-1} \\ &\Rightarrow x\alpha < x\mu^{-1} \\ &\Rightarrow x < x\mu^{-1}. \end{aligned}$$

Since $x \sim x\mu$, we get a contradiction. □

Lemma 6.3. *If $Nonsep(\kappa) \cap C_\alpha \neq \emptyset$, then the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: Let $x \in Nonsep(\kappa) \cap C_\alpha$. By Lemma 4.1, we may assume that $x \sim x\kappa$ but $x \neq x\kappa$. Set $\mathcal{T}_0 = \mathcal{T}_{\{x, x\kappa\}}$. We assume that $x < x\alpha$.

If $x\alpha^{-1} \in \mathcal{T}_0$ or $x\alpha \in \mathcal{T}_0$, then the ideal point $\hat{x} \in \widehat{\mathcal{T}}$ is fixed by κ and related to $\hat{x}\alpha$, and Lemma 4.1 applies.

So we may assume that $x\alpha, x\alpha^{-1} \notin \mathcal{T}_0$. Since $x < x\alpha$, either $\mathcal{T}_0 \subset x^+$ and $x\alpha \in y^-$ for some $y \sim x, y \neq x$ or $\mathcal{T}_0 \subset x^-$ and $x\alpha \in y^+$ for some $y \sim x, y \neq x$. We may assume the first possibility holds. Note that $\{x, y\} \subset [[x\alpha^{-1}, x\alpha]]$ and so $d(x, x\alpha) = 2n > 0$. In particular, we have $Nonsep(\alpha) = \emptyset$ and $C_\alpha = A_\alpha$ by Lemma 3.4 (Fig 18). We can also assume that $x \neq x\kappa^{p-18q}$ and $y \neq y\kappa^{p-18q}$ by Lemma 6.2.



FIGURE 18. C_α

Then we have 3 cases.

(1) $y\mu = y$ and $x \neq y\kappa^{p-18q}$.
$d(y, y\mu\alpha^2\mu) = d(y, y\alpha^2) = 4n$. But $d(y, y\alpha^3\kappa^{p-18q}\alpha^3) = d(y, y\kappa^{p-18q}\alpha^3) + d(y\kappa^{p-18q}\alpha^3, y\alpha^3\kappa^{p-18q}\alpha^3) = 6n + 6n = 12n$ (Fig 19). Since $n > 0$, this is impossible.


$y\alpha^3\kappa^{p-18q}$
$x\alpha^3\kappa^{p-18q}$
$y\kappa^{p-18q}$
$y\alpha^3\kappa^{p-18q}\alpha^3$
$x\alpha^3\kappa^{p-18q}\alpha^3$
$y\kappa^{p-18q}\alpha^3$
x
y
$x\alpha^3$
$y\alpha^3$

FIGURE 19. $x \neq y\kappa^{p-18q}$

(2) $y\mu = y$ and $x = y\kappa^{p-18q}$.

   (a) $y\beta \in A_\alpha \Leftrightarrow y\alpha^2\kappa^{p-18q} \in A_\alpha$.
   Then $d(x, y\alpha^2\kappa^{p-18q}) = d(y, y\alpha^2) = 4n$. Since $y\alpha^2 \in y^-$, $y\alpha^2\kappa^{p-18q} \in (y\kappa^{p-18q})^- = x^-$. Moreover $y\alpha^2\kappa^{p-18q} \in A_\alpha$. Therefore,

$$\begin{aligned} &y\alpha^2\kappa^{p-18q} = x\alpha^{-2} \\ &\Leftrightarrow y\beta = x\alpha^{-1} \\ &\Rightarrow y\beta < x \\ &\Leftrightarrow y\alpha^{-1} = y\mu^{-1}\alpha^{-1} = y\beta^2 < x\beta. \end{aligned}$$

   Since $x\beta \sim y\beta = x\alpha^{-1} \sim y\alpha^{-1}$, we get a contradiction.

   (b) $y\beta \notin A_\alpha$.
   Since $d(x\alpha, y\beta) = d(x\alpha, y\alpha^2\kappa^{p-18q}\alpha) = d(x, y\alpha^2\kappa^{p-18q}) = d(y, y\alpha^2) = 4n = d(x\alpha, x\alpha^{-1})$, $x\alpha^{-1} \notin [[x\alpha, y\beta]]$.
   If $x \in [[x\alpha, y\beta]]$, then $y\beta < x$ (Fig 20). Therefore $x\beta > y\alpha^{-1} = y\mu^{-1}\alpha^{-1} = y\beta^2$. Contradiction.
   If $x \notin [[x\alpha, x\beta]]$, $y > y\beta \Rightarrow y > y\beta^2 = y\alpha^{-1}$(Fig 20). Contradiction.



Figure 20. $x\beta$ and A_α

(3) $y \neq y\mu$.
   Since $x\mu\alpha^2 \sim y\alpha^2 \in y^-$, $x\mu\alpha^2 \in y^-$. Thus $x\mu\alpha^2\mu \in (y\mu)^- \subset y^+$. But $x\alpha^3\kappa^{p-18q}\alpha^3 \in y^- \cup \{y\}$(see Fig 19).

□

Lemma 6.4. *If $Nonsep(\kappa) \neq \emptyset$ and $Nonsep(\alpha) \cap C_\kappa \neq \emptyset$, then the action $G(p,q)$ on $\mathcal{T}$ is trivial.*

Proof: Let $x \in Nonsep(\alpha) \cap C_\kappa$. By Lemma 3.3, either $x \in Fix(\alpha)$ or x lies on some local axis $A^i_\kappa \approx \mathbb{R}$ (in $\mathcal{T}$) for κ. By Lemma 6.1, we may assume that x lies on some local axis A^i_κ. Then either $x \in Fix(\alpha)$ or the ideal point $\hat{x} \in \widehat{\mathcal{T}}$ is fixed by α and related to $\hat{x}\kappa$. In either case, Lemma 4.1 applies. □

Lemma 6.5. *If $G(p,q)$ acts nontrivially on $\mathcal{T}$, then:*

- *$C_\kappa \cup Nonsep(\kappa) \subset X_{j_0}$ for some $j_0 \in \mathcal{J}$, and*
- *$C_\alpha \cup Nonsep(\alpha) \subset T_{i_0}$ for some $i_0 \in \mathcal{I}$.*

Proof: By Lemma 6.4, $(C_\kappa \cup Nonsep(\kappa)) \cap Nonsep(\alpha) = \emptyset$. By Lemma 3.5 therefore, $C_\kappa \cup Nonsep(\kappa) \subset X_{j_0}$ for some $j_0 \in \mathcal{J}$. A symmetric argument proves the second statement. □

Proposition 6.6. *Suppose $Nonsep(\kappa) \neq \emptyset$. Then the action is trivial.*

Proof: Let i_0, j_0 be as guaranteed in Lemma 6.5. Suppose first that $T_{i_0}\kappa = T_{i_0}$. As remarked above, $A^{i_0}_\kappa \approx \mathbb{R}$. By Lemma 6.5, $Nonsep(\kappa) \cup A^{i_0}_\kappa \subset X_{j_0}$.

Consider first the possibility that $X_{j_0}\alpha = X_{j_0}$, and hence $A^{j_0}_{\alpha} \subset T_{i_0}$. In fact, $T_{i_0} \cap X_{j_0}$ is a subtree of $\mathcal{T}$ containing both $A^{i_0}_{\kappa}$ and $A^{j_0}_{\alpha}$. Therefore, if $A^{i_0}_{\kappa} \cap A^{j_0}_{\alpha} = \emptyset$, the bridge from $A^{i_0}_{\kappa}$ to $A^{j_0}_{\alpha}$ lies in $T_{i_0} \cap X_{j_0}$. If either of the two potential endpoints of $A^{i_0}_{\kappa}$(respectively, $A^{j_0}_{\alpha}$) exist in $\mathcal{T}$, they are in $Nonsep(\kappa)$ (respectively, $Nonsep(\alpha)$) and hence are not elements of $\mathcal{T}_{i_0}$ (respectively, X_{j_0}), and therefore cannot be on the bridge. Hence this bridge has the form $[[u, v]]$ or $[[u, v))$, where u and v are not separated from points in $A^{i_0}_{\kappa}$ and $A^{j_0}_{\alpha}$ respectively. Computing $A^{i_0}_{\kappa}$ in this case, we see that $A^{i_0}_{\kappa} \cap A^{i_0}_{\kappa}\alpha = \emptyset$, with the bridge from $A^{i_0}_{\kappa}$ to $A^{i_0}_{\kappa}\alpha$ of the form $[[u, w]]$ for some $w \sim w' \in A^{i_0}_{\kappa}\alpha$. So Lemma 5.5 reveals that the action of $G(p, q)$ on $\mathcal{T}$ is necessarily trivial. On the other hand, if $A^{i_0}_{\kappa} \cap A^{j_0}_{\alpha} \neq \emptyset$ then Lemma 6.1 guarantees that $A^{i_0}_{\kappa} \cap A^{j_0}_{\alpha} \subset [u, v]$ for some $u, v \in A^{i_0}_{\kappa}$. Computing $A^{i_0}_{\kappa}\alpha$ in this case, we see that one of the two conditions of Lemma 5.5 is satisfied, and so once again, the action of $G(p, q)$ on $\mathcal{T}$ must be trivial.

Next consider the possibility that $X_{j_0}\alpha = X_{j_1} \neq X_{j_0}$. Let y and $y\alpha$ denote the roots of X_{j_0} and $X_{j_0}\alpha$, respectively. Let $[[y, r]]$ denote the bridge from y to $A^{i_0}_{\kappa}$ in $\mathcal{T}$. By Lemma 6.1, we may assume that $r \sim r'$ for some $r' \in A^{i_0}_{\kappa}$. So $A^{i_0}_{\kappa} \cap A^{i_0}_{\kappa}\alpha = \emptyset$ with bridge $[[r, r\alpha]]$. Again, by Lemma 5.5, the action of $G(p, q)$ on $\mathcal{T}$ has a global fixed point.

Suppose that $T_{i_0}\kappa = T_{i_1} \neq T_{i_0}$. In particular, $x \neq x\kappa$. As shown in the proof of Proposition 8.7 in [RRS], we have $x\alpha \in x^-$ and $x \in (x\alpha)^-$ (Fig 21).



Figure 21. x and $x\alpha$ when $T_{i_0}\kappa = T_{i_1}$

If $x = x\mu = x\kappa^{p-18q}$, then $x = x\kappa^p$. Since $(p, q) = 1$, $x = x\kappa$. Hence x is not equal to at least one of $x\mu$ and $x\kappa^{p-18q}$.

Suppose that $x = x\alpha^2$. Assume first $x = x\kappa^{p-18q}$ (and hence $x \neq x\mu$). Then $x\alpha\mu^{-1} = x\alpha^3\mu^{-1} = x\alpha^{-3}\mu\alpha^2 = x\alpha\mu\alpha^2$. Since $d(x\mu^{-1}, x\alpha\mu^{-1}) = d(x, x\alpha) = d(x\mu, x\alpha\mu) = d(x\mu\alpha^2, x\alpha\mu\alpha^2)$ and $x\mu^{-1} \sim x = x\alpha^2 \sim x\mu\alpha^2$,

$$\begin{aligned}
&x\mu^{-1} = x\mu\alpha^2 \text{ (see Fig 22)}\\
&\Leftrightarrow x\alpha^2 = x = x\mu\alpha^2\mu = x\alpha\beta\alpha^2\\
&\Leftrightarrow x = x\alpha\beta\\
&\Leftrightarrow x\beta = x\alpha^2\kappa^{p-18q}\alpha = x\alpha = x\beta^{-1}\\
&\Rightarrow x\mu^{-1}\alpha^{-1} = x\beta^2 = x\\
&\Leftrightarrow x\mu^{-1} = x\alpha.
\end{aligned}$$

Since $x\mu^{-1} \sim x$, we get a contradiction. Now we can assume

Note that $x\alpha^3\kappa^{p-18q}\alpha^3 \in x^-$ (Fig 23). But $x\mu\alpha^2\mu \sim x\alpha^2\mu = x\mu \sim x$. So $x\mu\alpha^2\mu \in x^+ \cup \{x\}$. Contradiction.



Figure 22. $x = x\alpha^2 = x\kappa^{p-18q}$



FIGURE 23. Case (1) $x \neq x\kappa^{p-18q}, x \neq x\mu$



FIGURE 24. Case (2) $x \neq x\mu, x = x\kappa^{p-18q}$

Now we may assume $x \neq x\alpha^2$. Suppose that $x = x\alpha^3$. Then

$$x\alpha^3\kappa^{p-18q}\alpha^3 = x\kappa^{p-18q}\alpha^3 \sim x\alpha^3 = x.$$

But $x\mu\alpha^2\mu \not\sim x$ as shown in Fig 23(when $x \neq x\mu$) and Fig 25(when $x = x\mu$). Hence we also assume that $x \neq x\alpha^3$.

Recall that x is not equal to at least one of $x\mu$ and $x\kappa^{p-18q}$.

We have 3 cases. Note that $x\alpha^2, x\alpha^3 \in x^-$ and $x \in (x\alpha^2)^- \cap (x\alpha^3)^-$.

(1) $x \neq x\kappa^{p-18q}$ and $x \neq x\mu$.
As shown in Fig 23, $x\mu\alpha^2\mu \in x^+$ but $x\alpha^3\kappa^{p-18q}\alpha^3 \in x^-$.

(2) $x \neq x\mu$ and $x = x\kappa^{p-18q}$
It follows that $x\alpha^3\mu^{-1} = x\alpha^{-3}\mu\alpha^2$. Then $x\alpha^3\mu^{-1} \in x^+$, but $x\alpha^{-3}\mu\alpha^2 \in x^-$ (see Fig 24).

(3) $x \neq x\kappa^{p-18q}$ and $x = x\mu$.
As shown in the figure of case (1), $x\alpha^3\kappa^{p-18q}\alpha^3 \in (x\alpha^3)^+$. But $x\mu\alpha^2\mu = x\alpha^2\mu \in (x\alpha^3)^-$ (Fig 25).

□

REFERENCES

[BH] Bleiler, S., Hodgson, C., *Spherical space forms and Dehn surgery*, Topology, **35** (1996), no. 3, 809–833.

[BNR] Brittenham, M., Naimi, R., Roberts, R., *Graph manifolds and taut foliations*, J. Differential Geom., **45** (1997), no. 3, 446–470.

[CD] Calegari, D., Dunfield, N., *Laminations and groups of homeomorphisms of the circle*, Invent. Math., **152** (2003), no. 1, 149–204.

[CC] Candel, A., Conlon, L., *Foliations. II*, preprint.



FIGURE 25. Case (3) $x \neq x\kappa^{p-18q}, x = x\mu$

[Eu] Eudave-Muñoz, Mario, *Non-hyperbolic manifolds obtained by Dehn surgery on hyperbolic knots*, Geometric topology (Athens, GA, 1993), 35–61, AMS/IP Stud. Adv. Math., 2.1, Amer. Math. Soc., Providence, RI, 1997.

[FS] Fintushel, R., Stern, R., *Constructing lens spaces by surgery on knots*, Math. Z., **175** (1980), no. 1, 33–51.

[Ga] Gabai, David, *Problems in foliations and laminations*, Geometric Topology, AMS, Providence, RI, International Press, Cambridge, MA, 1997.

[GK] Gabai, D., Kazez, W., *Order trees and laminations of the plane*, Math. Res. Lett. **4** (1997), no. 4, 603–616.

[GO] Gabai, D.,Oertel, U., *Essential laminations in 3-manifolds*, Ann. of Math. (2) **130** (1989), no. 1, 41–73.

[HO] Hatcher, A., Oertel, U., *Boundary slopes for Montesinos knots*, Topology, **28** (1989), 453–480.

[Li] Li, Tao, *Laminar branched surfaces in 3-manifolds*, Geometry and Topology, Vol. 6 (2002), 153-194.

[No] Novikov, S., *Topology of foliations*, Trans. Moscow. Math. Soc., **14** (1965), 268–305.

[Oe] Oertel, Ulrich, *Closed incompressible surfaces in complements of star links*, Pacific J. Math., **111** (1984), 209–230.

[Pa] Palmeira, Frederico, *Open manifolds foliated by planes*, Ann. of Math., **107** (1978), 109–131.

[Rob] Roberts, Rachel, *Taut foliations in punctured surface bundles. I*, Proc. London Math. Soc., (3) 82 (2001), no. 3, 747–768.

[Ros] Rosenberg, Harold, *Foliations by planes*, Topology, **7** (1968), 131–138.

[Rol] Rolfsen, Dale, *Knots and links*, Mathematics Lecture Series, No. 7. Publish or Perish, Inc., Berkeley, Calif., 1976.

[RRS] Roberts, R., Shareshian, J., Stein, M., *Infinitely many hyperbolic 3-manifolds which contain no Reebless foliation.*, J. Amer. Math. Soc., **16** (2003), 639–679.

[We] Weeks, Jeffrey, SNAPPEA, the hyperbolic structures computer program.

DEPARTMENT OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 151-747, KOREA
E-mail address: `jhjun@math.snu.ac.kr`